

LA-UR-21-22464

Approved for public release; distribution is unlimited.

Title: Neutrinos and the Dark Sector

Author(s): Louis, William Charles III

Intended for: Present at the P-Division Cafe

Issued: 2021-03-12

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Neutrinos & the Dark Sector

W. C. Louis, LANL

January 21, 2021

- Neutrino Properties
- Short Baseline Neutrino Anomalies
- Dark Sector Models
- SBN Project at Fermilab
- CCM Experiment at Lujan
- Conclusion

Standard Model of Elementary Particles

three generations of matter (fermions)			interactions / force carriers (bosons)	
	I	II	III	
QUARKS	$\approx 2.2 \text{ MeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$ u up	$\approx 1.28 \text{ GeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$ c charm	$\approx 173.1 \text{ GeV}/c^2$ $\frac{2}{3}$ $\frac{1}{2}$ t top	$\approx 124.97 \text{ GeV}/c^2$ 0 0 1 g gluon
	$\approx 4.7 \text{ MeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ d down	$\approx 96 \text{ MeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ s strange	$\approx 4.18 \text{ GeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ b bottom	0 0 1 γ photon
	$\approx 0.511 \text{ MeV}/c^2$ -1 $\frac{1}{2}$ e electron	$\approx 105.66 \text{ MeV}/c^2$ -1 $\frac{1}{2}$ μ muon	$\approx 1.7768 \text{ GeV}/c^2$ -1 $\frac{1}{2}$ τ tau	$\approx 91.19 \text{ GeV}/c^2$ 0 1 Z Z boson
LEPTONS	$< 2.2 \text{ eV}/c^2$ 0 $\frac{1}{2}$ ν_e electron neutrino	$< 0.17 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ ν_μ muon neutrino	$< 18.2 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ ν_τ tau neutrino	$\approx 80.39 \text{ GeV}/c^2$ ± 1 1 W W boson
				SCALAR BOSONS GAUGE BOSONS VECTOR BOSONS

The Standard Model makes up only $\sim 5\%$ of the mass-energy of the universe!

The dark sector (dark matter & dark energy) makes up $\sim 95\%$ of the mass-energy of the universe!

Are neutrinos the portal to the dark sector?

Salient Features of Neutrinos


- Neutrinos are neutral, spin $\frac{1}{2}$, point-like particles with small, non-zero masses ($< \sim 1$ eV)
- Neutrinos interact very weakly with matter
- Neutrinos & photons dominate the universe in terms of number of particles
- Neutrinos undergo oscillations

Neutrino Oscillations

Weak Eigenstates

Eigenstates of Propagation


$$\begin{aligned} \nu_\mu &= \cos\theta \nu_1 + \sin\theta \nu_2 \\ \nu_e &= -\sin\theta \nu_1 + \cos\theta \nu_2 \end{aligned}$$



$$P_{\nu_\mu \rightarrow \nu_e} = \sin^2(2\theta) \sin^2(1.27 \Delta m^2 L/E_\nu)$$

$$\Delta m^2 = m_2^2 - m_1^2 \text{ in eV}^2, L \text{ in meters, } E_\nu \text{ in MeV}$$



Observation of ν Oscillations



SuperK, SNO, KamLAND, BOREXINO

$$\Delta m^2 \sim 0.00007 \text{ eV}^2$$


Other sources of neutrinos:



SuperK, K2K, MINOS, OPERA, T2K, Double Chooz, Daya Bay, RENO, NOvA

$$\Delta m^2 \sim 0.002 \text{ eV}^2$$

2015 Nobel Prize
for SuperK &
SNO!




LSND, MiniBooNE, Reactor ν , Gallium Anomaly

$$\Delta m^2 \sim 1 \text{ eV}^2$$


Short-Baseline Neutrino Anomalies

$\nu_\mu \rightarrow \nu_e$ Appearance


LSND




New MiniBooNE Neutrino




New MiniBooNE Neutrino & Antineutrino




LSND



New MiniBooNE Neutrino




New MiniBooNE Neutrino + Antineutrino




Short-Baseline Neutrino Anomalies

ν_μ & ν_e Disappearance


Gallium Anomaly




Neutrino-4 Reactor Experiment




Reactor Neutrino Anomaly



IceCube Experiment



3+N Sterile Neutrino Models



3+N Models with Dark Sector Particles Can Explain the World Data

(Sterile neutrinos with new mediators, interactions, and scalar particles!)

- Sterile Neutrino Decay
- Sterile Neutrinos with Resonant Oscillations
- Sterile Neutrino NSI & New Gauge Bosons
- Light WIMP Production (Light WIMPs can behave like neutrinos)
- Lorentz Violation & CPT Violation
- Extra Dimensions (active neutrinos are stuck on the brane, while sterile neutrinos can propagate in the bulk)
- Mass-Varying Neutrinos
- Neutrino Decoherence
- etc.

A two-Higgs doublet solution to the LSND, MiniBooNE and muon $g - 2$ anomalies

Waleed Abdallah, Raj Gandhi, and Samiran Roy

arXiv:2010.06159








FIG. 1: Feynman diagram of the scattering process in our model which leads to the excess in LSND and MB.

A two-Higgs doublet solution to the LSND, MiniBooNE and muon $g - 2$ anomalies

Waleed Abdallah, Raj Gandhi, and Samiran Roy


arXiv:2010.06159



Neutrinos & the Dark Sector

(Daniele Alves & Patrick deNiverville)

- Recent theoretical effort initiated at LANL: implications of dark sectors to neutrino phenomenology beyond the Standard Model




- Under generic assumptions above, neutrino oscillations in medium should be modified by *new matter effects*
(D.S.M. Alves, P. deNiverville, W. Louis, *in preparation*).

Neutrinos & the Dark Sector

(Daniele Alves & Patrick deNiverville)



- Oscillation probabilities can be enhanced for specific windows of neutrino energy and medium density

- could provide explanation for excess of low energy electron-neutrinos observed by MiniBooNE
- does not affect of MSW matter effect in the Sun



Short Baseline Neutrino (SBN) Program


See talk by D. Schmitz



ICARUS-T600

M. Touns


476 tons



MicroBooNE

First Results From MicroBooNE

89 tons



Short Baseline Near Detector (SBND)


112 tons

5


CCM Experiment at Lujan

Lujan Target

- Intense protons (100 μA)
- Fast Beam (295 nsec)
- Compact source



Intense source muon neutrinos: target MCNP simulation flux $4.74 \times 10^5 \text{ nu/cm}^2/\text{s}$ at 20 m (slide 52)



Lujan Experimental Area

- Space for large 10-ton liquid Argon neutrino detector.
- Run detector in multiple locations.
- Room to deploy shielding, large overhead crane, power, etc

Conclusion


- The results from LSND & MiniBooNE and the other anomalies in short baseline ν experiments cannot be explained by the 3 ν paradigm and suggest the existence of sterile neutrinos and new dark sector particles and interactions.
- Sterile neutrinos & the new dark sector particles would contribute to the dark matter of the universe and would have a big impact on particle physics, nuclear physics, astrophysics and cosmology.
- Future experiments (e.g. SBN at Fermilab & CCM at Lujan) have the golden opportunity of proving whether sterile neutrinos and dark sector particles exist!

Backup

Coherent CAPTAIN-Mills (CCM) experiment

Production mechanism:


charged pions decaying at rest \Rightarrow monoenergetic neutrinos




Detection mechanism:

Coherent Elastic Neutrino-Nucleus Scattering
“CEvNS”



present for all active neutrinos



absent for sterile neutrinos




Detecting Coherent Neutrinos: Maximizing Scintillation Light Detection!



- Liquid Argon scintillates at 128 nm with 40 photon/keV
 - fast 6 nsec and slower ~1 usec time constants.
- Detailed RAT/GEANT4 simulation predicts 10-20 keV detection threshold.
- CCM will also allow more tests of LAr scintillation light for SBN & DUNE.

CCM Sensitivity

$\nu_\mu \rightarrow \nu_e$ Appearance



Reaction	L = 20 m (events/yr)	L = 40 m (events/yr)
Coherent ν_μ (E = 30 MeV)	2709	677
Coherent $\nu_e + \bar{\nu}_\mu$	9482	2370
Charged Current ν_e	257	64
Neutral Current ν_μ	36	18
Neutral Current $\bar{\nu}_\mu$	79	20

- Can prove/disprove at $\sim 3\sigma$ LSND 3+1 sterile neutrino hypothesis.
- Five year run would approach 5σ !
- If no signal, can rule out world best fit at better than 90%

Future Short-Baseline ν Experiments

- There is a diverse set of experiments, spanning vastly different energy Scales (from ~ 1 MeV to ~ 10 TeV), that have been proposed to test the 3+N models & resolve the present anomalies:

- Accelerator ν Experiments: **MicroBooNE+SBND+ICARUS**, MINOS+, NOvA, **DUNE**, OscSNS at ORNL, **CAPTAIN-Mills**, JSNS² (J-PARC E56), IsoDAR, nuPRISM






Figure 7. The ICARUS T600 detector installed in Hall B at LNGS.

- Reactor ν Experiments: Neutrino-4 SOLID, PROSPECT, NEOS, DANSS





- Radioactive Source ν Experiments: BEST





- Atmospheric ν Experiments: IceCube




- Beta Decay & Double Beta Decay



ν_e Appearance Sensitivity



ν_μ Disappearance Sensitivity



Sensitivity includes full flux and cross section systematics,
but not detector systematics at this time.

Probability of Neutrino Oscillations

$$P_{\alpha\beta} = \delta_{\alpha\beta} - 4 \sum_i \sum_j |U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j}| \sin^2(1.27 \Delta m_{ij}^2 L/E_\nu)$$

As #v increases, the formalism gets rapidly more complicated!

#v	# Δm_{ij}^2	# θ_{ij}	#CP Phases
2	1	1	0
3	2	3	1
4	3	6	3
5	4	10	6
6	5	15	10

Therefore, there needs to be ≥ 3 neutrino mixing for CP Violation!

3+N Models With ν_e Appearance Require Large ν_e & ν_μ Disappearance!

In general, $P(\nu_\mu \rightarrow \nu_e) \sim \frac{1}{4} P(\nu_\mu \rightarrow \nu_x) P(\nu_e \rightarrow \nu_x)$

Assuming that the 3 light neutrinos are mostly active and the N heavy neutrinos are mostly sterile.

For 3+1 Models:
arXiv:1207.4765

$$P(\nu_\alpha \rightarrow \nu_\beta) \simeq 4|U_{\alpha 4}|^2|U_{\beta 4}|^2 \sin^2(1.27\Delta m_{41}^2 L/E) ,$$

$$P(\nu_\alpha \rightarrow \nu_\alpha) \simeq 1 - 4(1 - |U_{\alpha 4}|^2)|U_{\alpha 4}|^2 \sin^2(1.27\Delta m_{41}^2 L/E) .$$

3+N Models With ν_e Appearance Require Large ν_e & ν_μ Disappearance!

In general, $P(\nu_\mu \rightarrow \nu_e) \sim \frac{1}{4} P(\nu_\mu \rightarrow \nu_x) P(\nu_e \rightarrow \nu_x)$

Assuming that the 3 light neutrinos are mostly active
and the N heavy neutrinos are mostly sterile.

For 3+2 Models:
arXiv:1207.4765

$$\begin{aligned}
 P(\nu_\alpha \rightarrow \nu_\beta) \simeq & -4|U_{\alpha 5}||U_{\beta 5}||U_{\alpha 4}||U_{\beta 4}| \cos \phi_{54} \sin^2(1.27\Delta m_{54}^2 L/E) \\
 & +4(|U_{\alpha 4}||U_{\beta 4}| + |U_{\alpha 5}||U_{\beta 5}| \cos \phi_{54})|U_{\alpha 4}||U_{\beta 4}| \sin^2(1.27\Delta m_{41}^2 L/E) \\
 & +4(|U_{\alpha 4}||U_{\beta 4}| \cos \phi_{54} + |U_{\alpha 5}||U_{\beta 5}|)|U_{\alpha 5}||U_{\beta 5}| \sin^2(1.27\Delta m_{51}^2 L/E) \\
 & +2|U_{\beta 5}||U_{\alpha 5}||U_{\beta 4}||U_{\alpha 4}| \sin \phi_{54} \sin(2.53\Delta m_{54}^2 L/E) \\
 & +2(|U_{\alpha 5}||U_{\beta 5}| \sin \phi_{54})|U_{\alpha 4}||U_{\beta 4}| \sin(2.53\Delta m_{41}^2 L/E) \\
 & +2(-|U_{\alpha 4}||U_{\beta 4}| \sin \phi_{54})|U_{\alpha 5}||U_{\beta 5}| \sin(2.53\Delta m_{51}^2 L/E) ,
 \end{aligned}$$

ϕ_{54} is the CP
Phase angle

$$\begin{aligned}
 P(\nu_\alpha \rightarrow \nu_\alpha) \simeq & 1 - 4|U_{\alpha 4}|^2|U_{\alpha 5}|^2 \sin^2(1.27\Delta m_{54}^2 L/E) \\
 & -4(1 - |U_{\alpha 4}|^2 - |U_{\alpha 5}|^2)(|U_{\alpha 4}|^2 \sin^2(1.27\Delta m_{41}^2 L/E) \\
 & + |U_{\alpha 5}|^2 \sin^2(1.27\Delta m_{51}^2 L/E)) .
 \end{aligned}$$